

Pediatric Cardiac Arrest Algorithm




American Heart Association

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



Pediatric Advanced Life Support



CPR Quality

- Push hard ($\geq 1/3$ of anteroposterior diameter of chest) and fast (100-120/min) and allow complete chest recoil
- Minimize interruptions in compressions
- Change compressor every 2 minutes, or sooner if fatigued
- If no advanced airway, 15:2 compression-ventilation ratio
- If advanced airway, provide continuous compressions and give a breath every 2-3 seconds

Shock Energy for Defibrillation

- First shock 2 J/kg
- Second shock 4 J/kg
- Subsequent shocks ≥ 4 J/kg, maximum 10 J/kg or adult dose

Drug Therapy

- **Epinephrine IV/IO dose:** 0.01 mg/kg (0.1 mL/kg of the 0.1 mg/mL concentration). Max dose 1 mg. Repeat every 3-5 minutes. If no IV/IO access, may give endotracheal dose: 0.1 mg/kg (0.1 mL/kg of the 1 mg/mL concentration).
- **Amiodarone IV/IO dose:** 5 mg/kg bolus during cardiac arrest. May repeat up to 3 total doses for refractory VF/pulseless VT or
- **Lidocaine IV/IO dose:** Initial: 1 mg/kg loading dose

Advanced Airway

- Endotracheal intubation or supraglottic advanced airway
- Waveform capnography or capnometry to confirm and monitor ET tube placement

Reversible Causes

- Hypovolemia
- Hypoxia
- Hydrogen ion (acidosis)
- Hypoglycemia
- Hypo-/hyperkalemia
- Hypothermia
- Tension pneumothorax
- Tamponade, cardiac
- Toxins
- Thrombosis, pulmonary
- Thrombosis, coronary



Pediatric Bradycardia With a Pulse Algorithm




American Heart Association.

American Academy of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

Pediatric Advanced Life Support



Doses/Details

Epinephrine IV/IO dose:

0.01 mg/kg (0.1 mL/kg of the 0.1 mg/mL concentration). Repeat every 3-5 minutes. If IV/IO access not available but endotracheal (ET) tube in place, may give ET dose: 0.1 mg/kg (0.1 mL/kg of the 1 mg/mL concentration).

Atropine IV/IO dose:

0.02 mg/kg. May repeat once. Minimum dose 0.1 mg and maximum single dose 0.5 mg.

Possible Causes

- Hypothermia
- Hypoxia
- Medications

Pediatric Tachycardia With a Pulse Algorithm




American Heart Association.

American Academy of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

Pediatric Advanced Life Support



Pediatric Septic Shock Algorithm




American Heart Association.

American Academy of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

Pediatric Advanced Life Support



Brierley J, Carcillo JA, Choong K, et al. Clinical practice parameters for hemodynamic support of pediatric and neonatal septic shock: 2007 update from the American College of Critical Care Medicine. *Crit Care Med*. 2009;37(2):666-688. Kissoon N, Orr RA, Carcillo JA. Updated American College of Critical Care Medicine—pediatric advanced life support guidelines for management of pediatric and neonatal septic shock: relevance to the emergency care clinician. *Pediatr Emerg Care*. 2010;26(11):867-869.

Management of Shock After ROSC Algorithm




American
Heart
Association.

American Academy
of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

Pediatric Advanced Life Support



Estimation of Maintenance Fluid Requirements

- **Infants <10 kg:** 4 mL/kg per hour

Example: For an 8-kg infant, estimated maintenance fluid rate
 $= 4 \text{ mL/kg per hour} \times 8 \text{ kg}$
 $= 32 \text{ mL per hour}$

- **Children 10-20 kg:** 4 mL/kg per hour for the first 10 kg + 2 mL/kg per hour for each kg above 10 kg

Example: For a 15-kg child, estimated maintenance fluid rate
 $= (4 \text{ mL/kg per hour} \times 10 \text{ kg})$
 $+ (2 \text{ mL/kg per hour} \times 5 \text{ kg})$
 $= 40 \text{ mL/hour} + 10 \text{ mL/hour}$
 $= 50 \text{ mL/hour}$

- **Children >20 kg:** 4 mL/kg per hour for the first 10 kg + 2 mL/kg per hour for 11-20 kg + 1 mL/kg per hour for each kg above 20 kg.

Example: For a 28-kg child, estimated maintenance fluid rate
 $= (4 \text{ mL/kg per hour} \times 10 \text{ kg})$
 $+ (2 \text{ mL/kg per hour} \times 10 \text{ kg})$
 $+ (1 \text{ mL/kg per hour} \times 8 \text{ kg})$
 $= 40 \text{ mL per hour} + 20 \text{ mL per hour}$
 $+ 8 \text{ mL per hour}$
 $= 68 \text{ mL per hour}$

After initial stabilization, adjust the rate and composition of intravenous fluids based on the patient's clinical condition and state of hydration. In general, provide a continuous infusion of a dextrose-containing solution for infants. Avoid hypotonic solutions in critically ill children; for most patients use isotonic fluid such as normal saline (0.9% NaCl) or lactated Ringer's solution with or without dextrose, based on the child's clinical status.

*Milrinone can cause hypotension, so use and initiation of it should generally be reserved for those experienced with its use, initiation, and side effects (eg, ICU personnel).



Pediatric Color-Coded Length-Based Resuscitation Tape



American
Heart
Association.

American Academy
of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



Pediatric Advanced Life Support

Zone	3 kg	4 kg	5 kg	Pink	Red	Purple	Yellow	White	Blue	Orange	Green
ETT uncuffed (mm)	3.5	3.5	3.5	3.5	3.5	4.0	4.5	5.0	5.5	N/A	N/A
ETT cuffed (mm)	3.0	3.0	3.0	3.0	3.0	3.5	4.0	4.5	5.0	5.5	6.0
Lip-tip (cm)	9-9.5	9.5-10	10-10.5	10-10.5	10.5-11	11-12	12.5-13.5	14-15	15.5-16.5	17-18	18.5-19.5
Suction (F)	8	8	8	8	8	8	10	10	10	10	12
L-scope blade	1 straight	1 straight	1 straight	1 straight	1 straight	1-1.5 straight	2 straight/ curved	2 straight/ curved	2 straight/ curved	2-3 straight/ curved	2-3 straight/ curved
Stylet	6 F	6 F	6 F	6 F	6 F	6 F	10 F	10 F	10 F	14 F	14 F
OPA (mm)	50	50	50	50	50	60	60	60	70	80	80
NPA (F)	14	14	14	14	14	18	20	22	24	26	26
Bag-mask device (minimum mL)	450	450	450	450	450	450	450	450-750	750-1000	750-1000	1000
ETCO ₂ detector	Ped	Ped	Ped	Ped	Ped	Ped	Ped	Adult	Adult	Adult	Adult
LMA	1	1	1	1.5	1.5	2	2	2	2-2.5	2.5	3
Tidal volume (mL)	20-30	24-40	30-50	40-65	50-85	65-105	80-130	100-165	125-210	160-265	200-330
Frequency	20-25/min	20-25/min	20-25/min	20-25/min	20-25/min	15-25/min	15-25/min	15-25/min	12-20/min	12-20/min	12-20/min

Abbreviations: ETT, endotracheal tube; F, French; LMA, laryngeal mask airway; NPA, nasopharyngeal airway; OPA, oropharyngeal airway; Ped, pediatric.

The Broselow-Luten System Point of Care Guide is © 2020 Vyair Medical, Inc.; used with permission.

PALS Systematic Approach Algorithm




American Heart Association.

American Academy of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

Pediatric Advanced Life Support



Components of Post-Cardiac Arrest Care



American
Heart
Association.

American Academy
of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

Pediatric Advanced Life Support

Oxygenation and ventilation	Check
Measure oxygenation and target normoxemia 94%-99% (or child's normal/appropriate oxygen saturation).	<input type="checkbox"/>
Measure and target $Paco_2$ appropriate to the patient's underlying condition and limit exposure to severe hypercapnia or hypocapnia.	<input type="checkbox"/>
Hemodynamic monitoring	
Set specific hemodynamic goals during post-cardiac arrest care and review daily.	<input type="checkbox"/>
Monitor with cardiac telemetry.	<input type="checkbox"/>
Monitor arterial blood pressure.	<input type="checkbox"/>
Monitor serum lactate, urine output, and central venous oxygen saturation to help guide therapies.	<input type="checkbox"/>
Use parenteral fluid bolus with or without inotropes or vasopressors to maintain a systolic blood pressure greater than the fifth percentile for age and sex.	<input type="checkbox"/>
Targeted temperature management (TTM)	
Measure and continuously monitor core temperature.	<input type="checkbox"/>
Prevent and treat fever immediately after arrest and during rewarming.	<input type="checkbox"/>
If patient is comatose apply TTM (32°C-34°C) followed by (36°C-37.5°C) or only TTM (36°C-37.5°C).	<input type="checkbox"/>
Prevent shivering.	<input type="checkbox"/>
Monitor blood pressure and treat hypotension during rewarming.	<input type="checkbox"/>
Neuromonitoring	
If patient has encephalopathy and resources are available, monitor with continuous electroencephalogram.	<input type="checkbox"/>
Treat seizures.	<input type="checkbox"/>
Consider early brain imaging to diagnose treatable causes of cardiac arrest.	<input type="checkbox"/>
Electrolytes and glucose	
Measure blood glucose and avoid hypoglycemia.	<input type="checkbox"/>
Maintain electrolytes within normal ranges to avoid possible life-threatening arrhythmias.	<input type="checkbox"/>
Sedation	
Treat with sedatives and anxiolytics.	<input type="checkbox"/>
Prognosis	
Always consider multiple modalities (clinical and other) over any single predictive factor.	<input type="checkbox"/>
Remember that assessments may be modified by TTM or induced hypothermia.	<input type="checkbox"/>
Consider electroencephalogram in conjunction with other factors within the first 7 days after cardiac arrest.	<input type="checkbox"/>
Consider neuroimaging such as magnetic resonance imaging during the first 7 days.	<input type="checkbox"/>